arXiv:1604.02865v1 [math.SP] 11 Apr 2016

Spectrum and genus of commuting graphs of some classes of finite rings

Jutirekha Dutta and Rajat Kanti Nath*

Department of Mathematical Sciences,
Tezpur University, Napaam-784028, Sonitpur, Assam, India.
Emails: jutirekhadutta@yahoo.com, rajatkantinath@yahoo.com

Abstract: The commuting graph of a non-commutative ring R with center $Z(R)$ is a simple undirected graph whose vertex set is $R \setminus Z(R)$ and two vertices x, y are adjacent if and only if $xy = yx$. In this paper, we compute the spectrum and genus of commuting graphs of some classes of finite rings.

Key words: Integral graph, Commuting graph, Spectrum of graph.
2010 Mathematics Subject Classification: 05C25, 05C50, 16P10

1 Introduction

Let R be a non-commutative ring with center $Z(R)$. The commuting graph of R, denoted by Γ_R, is a simple undirected graph whose vertex set is $R \setminus Z(R)$ and two vertices x, y are adjacent if and only if $xy = yx$. In recent years, many mathematicians have considered commuting graph of different rings and studied various graph theoretic aspects (see [1, 4, 5, 10, 14, 16, 17]). Some generalizations of Γ_R are also considered in [3, 8]. In this paper, we compute the spectrum and genus of the commuting graph of some classes of finite rings.

Recall that the spectrum of a graph $\mathcal{G}$ denoted by $\mathrm{Spec}(\mathcal{G})$ is the set $\{\lambda_1^{k_1}, \lambda_2^{k_2}, \ldots, \lambda_n^{k_n}\}$, where $\lambda_1, \lambda_2, \ldots, \lambda_n$ are the eigenvalues of the adjacency matrix of $\mathcal{G}$ with multiplicities $k_1, k_2, \ldots, k_n$ respectively. A graph $\mathcal{G}$ is called integral if $\mathrm{Spec}(\mathcal{G})$ contains only integers. It is well-known that the complete graph K_n on n vertices

*Corresponding author

is integral and $\mathrm{Spec}(K_n) = \{(-1)^{n-1}, (n-1)^1\}$. Further, if $\mathcal{G} = \bigsqcup_{i=1}^{l} K_{m_i}$, where K_{m_i} are complete graphs on m_i vertices for $1 \leq i \leq l$, then

$$\mathrm{Spec}(\mathcal{G}) = \{(-1)^{\sum_{i=1}^{l} m_i - l}, (m_1 - 1)^1, (m_2 - 1)^1, \ldots, (m_l - 1)^1\}. \tag{1.1}$$

The notion of integral graph was introduced by Harary and Schwenk [11] in the year 1974. Since then many mathematicians have considered integral graphs, see for example [2, 12, 18]. In [9, 15], the authors have determined several groups whose commuting graphs are integral.

The genus of a graph is the smallest non-negative integer n such that the graph can be embedded on the surface obtained by attaching n handles to a sphere. We write $\gamma(\mathcal{G})$ to denote the genus of a graph $\mathcal{G}$. It is worth mentioning that

$$\gamma(K_n) = \left\lceil \frac{(n-3)(n-4)}{12} \right\rceil \text{ if } n \geq 3 \text{ (see [19, Theorem 6-38])}.$$

Also, if $\mathcal{G} = \bigsqcup_{B \in \mathcal{B}} B$, where $\mathcal{B}$ denotes the block decomposition of $\mathcal{G}$, then by [6, Corollary 1] we have

$$\gamma(\mathcal{G}) = \sum_{B \in \mathcal{B}} \gamma(B). \tag{1.2}$$

A graph $\mathcal{G}$ is called planar or toroidal if $\gamma(\mathcal{G}) = 0$ or 1 respectively. As a consequence of our results we show that the commuting graphs of non-commutative rings of order p^2 and p^3 are integral but not toroidal.

For any element r of a ring R, the set $C_R(r) = \{s \in R : rs = sr\}$ is called the centralizer of r in R. Let $|\operatorname{Cent}(R)| = |\{C_R(r) : r \in R\}|$, that is the number of distinct centralizers in R. A ring R is called an n-centralizer ring if $|\operatorname{Cent}(R)| = n$. In [7], Dutta et al. have characterized finite n-centralizer rings for $n = 4, 5$. As a consequence of our results, we show that the commuting graphs of $4, 5$-centralizer finite rings are integral but not toroidal. Further, we show that the commuting graph of a finite $(p+2)$-centralizer p-ring is integral for any prime p. We conclude this paper by computing the spectrum and genus of the commuting graphs of some finite rings with some specific commuting probability.

2 Main results

A non-commutative ring R is called a CC-ring if all the centralizers of its non-central elements are commutative. In [10], Erfanian et al. have initiated the study of CC-rings. In particular, they have computed the diameter of Γ_R^c and showed that the clique number and chromatic number of Γ_R^c are same for a CC-ring R,

where Γ_R^c denotes the complement of Γ_R. In the following theorem we compute the spectrum and genus of Γ_R for a finite CC-ring R.

Theorem 2.1. *Let R be a finite CC-ring with distinct centralizers $S_1, S_2, \ldots, S_n$ of non-central elements of R. Then*

$$\mathrm{Spec}(\Gamma_R) = \{(-1)^{\sum_{i=1}^{n}|S_i|-n(|Z(R)|+1)}, (|S_1|-|Z(R)|-1)^1, \ldots, (|S_n|-|Z(R)|-1)^1\}$$

and $\gamma(\Gamma_R) = \sum_{i=1}^{n}\gamma(K_{|S_i|-|Z(R)|})$.

Proof. Let R be a finite CC-ring and $S_1, S_2, \ldots, S_n$ be the distinct centralizers of non-central elements of R. Let $S_i = C_R(s_i)$ where $s_i \in R \setminus Z(R)$ for $i = 1, 2, \ldots, n$. Let $s \in (S_i \cap S_j) \setminus Z(R)$ for some i, j such that $1 \leq i \neq j \leq n$. Then s commutes with s_i as well as s_j. Let $t \in C_R(s)$ then $ts_i = s_it$ since $s_i \in C_R(s)$ and R is a CC-ring. Therefore $t \in C_R(s_i)$ and so $C_R(s) \subseteq C_R(s_i)$. Again, let $u \in C_R(s_i)$ then $us = su$ since $s \in S_i = C_R(s_i)$ and R is a CC-ring. Therefore, $u \in C_R(s)$ and so $C_R(s_i) \subseteq C_R(s)$. Thus $C_R(s) = C_R(s_i)$. Similarly, it can be seen that $C_R(s) = C_R(s_j)$. Hence $C_R(s) = C_R(s_i) = C_R(s_j)$ which is a contradiction. Therefore, $S_i \cap S_j = Z(R)$ for $1 \leq i \neq j \leq n$. This shows that $\Gamma_R = \bigsqcup_{i=1}^{n} K_{|S_i|-|Z(R)|}$. Now the results follow from (1.1) and (1.2). □

Corollary 2.2. *Let R be a finite CC-ring and A be any finite commutative ring. Then the spectrum of the commuting graph of $R \times A$ is given by*

$$\{(-1)^{\sum_{i=1}^{n}|A|(|S_i|-|Z(R)|)-n}, (|A|(|S_1|-|Z(R)|)-1))^1, \ldots, (|A|(|S_n|-|Z(R)|)-1))^1\}$$

and $\gamma(\Gamma_R) = \sum_{i=1}^{n}\gamma(K_{|A|(|S_i|-|Z(R)|)})$ *where $S_1, \ldots, S_n$ are the distinct centralizers of non-central elements of R.*

Proof. Note that $Z(R \times A) = Z(R) \times A$ and $S_1 \times A, S_2 \times A, \ldots, S_n \times A$ are the distinct centralizers of non-central elements of $R \times A$. Therefore, if R is an CC-ring then $R \times A$ is also a CC-ring. Hence, the result follows from Theorem 2.1. □

In general it is difficult to determine all the finite non-commutative rings whose commuting graphs are integral. However, by Theorem 2.1, it follows that the commuting graph of a finite CC-ring is integral. Further, if R is a finite CC-ring and A is any finite commutative ring then, by Corollary 2.2, the commuting graph of $R \times A$ is also integral. In the next result we consider a particular class of CC-rings and compute the spectrum and genus of its commuting graph.

Theorem 2.3. *Let R be a finite ring such that the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime. Then*

$$\mathrm{Spec}(\Gamma_R) = \{(-1)^{(p^2-1)|Z(R)|-p-1}, ((p-1)|Z(R)|-1)^{p+1}\} \text{ and}$$

$$\gamma(\Gamma_R) = (p+1)\gamma(K_{(p-1)|Z(R)|}).$$

Proof. Let $\frac{R}{Z(R)} = \langle a + Z(R), b + Z(R) : pa, pb \in Z(R)\rangle$, where $a, b \in R$ with $ab \neq ba$. Then for any $z \in Z(R)$, we have

$$\begin{aligned} C_R(a) &= C_R(ia + z) \\ &= Z(R) \sqcup a + Z(R) \sqcup \cdots \sqcup (p-1)a + Z(R) \text{ for } 1 \leq i \leq p-1; \end{aligned}$$

and for $1 \leq j \leq p$,

$$\begin{aligned} C_R(ja + b) &= C_R(ja + b + z) \\ &= Z(R) \sqcup (ja + b) + Z(R) \sqcup \cdots \sqcup ((p-1)ja + (p-1)b) + Z(R). \end{aligned}$$

These are the only centralizers of non-central elements of R. Also note that these centralizers are commutative subrings of R. Thus R is a CC-ring and

$$\Gamma_R = K_{|C_R(a)\setminus Z(R)|} \sqcup (\bigsqcup_{j=1}^{p} K_{|C_R(ja+b)\setminus Z(R)|}).$$

That is, $\Gamma_R = K_{(p-1)|Z(R)|} \sqcup (\bigsqcup_{j=1}^{p} K_{(p-1)|Z(R)|}) = \Gamma_R = \bigsqcup_{j=1}^{p+1} K_{(p-1)|Z(R)}$, since $|C_R(a)| = p|Z(R)|$ and $|C_R(ja + b)| = p|Z(R)|$ for $1 \leq j \leq p$. Hence the results follow from (1.1) and (1.2). □

3 Some consequences

In this section, we obtain several consequences of the results obtained in Section 2. In general it is difficult to determine all finite non-commutative rings whose commuting graphs are planar or toridal. In this section, we characterize some finite rings whose commuting graphs are planar. We begin with the following result.

Proposition 3.1. *Let R be a finite ring such that the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$, where p is a prime. Then*

(a) *Γ_R is integral but not toroidal.*

(b) Γ_R *is planar if and only if* $p = 2, |Z(R)| = 1, 2, 3$ *or* 4*; or* $p = 3, |Z(R)| = 1$ *or* 2.

Proof. Part (a) follows from Theorem 2.3. If $p = 2$ then $\gamma(\Gamma_R) = 0$ if and only if $3\gamma(K_{|Z(R)|}) = 0$ if and only if $|Z(R)| = 1, 2, 3$ or 4. If $p = 3$ then $\gamma(\Gamma_R) = 0$ if and only if $4\gamma(K_{2|Z(R)|}) = 0$ if and only if $|Z(R)| = 1$ or 2. Hence, part (b) follows. □

Proposition 3.2. *Let* p *be a prime and* R *a non-commutative ring of order* p^2*. Then*

(a) Γ_R *is integral but not toroidal.*

(b) Γ_R *is planar if and only if* $p = 2, 3$ *or* 5.

Proof. Note that $|Z(R)| = 1$ and the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. So, by Theorem 2.3, we have

$$\text{Spec}(\Gamma_R) = \{(-1)^{p^2-p-2}, (p-2)^{p+1}\} \text{ and } \gamma(\Gamma_R) = (p+1)\gamma(K_{p-1}).$$

Thus it follows that Γ_R is integral. Also, $\gamma(\Gamma_R) \neq 1$, that is, Γ_R is not toroidal. Part (b) follows from the fact that $\gamma(\Gamma_R) = 0$ if and only if $p = 2, 3$ or 5. □

Proposition 3.3. *Let* p *be a prime and* R *a non-commutative ring of order* p^3*. Then*

(a) Γ_R *is integral but not toroidal.*

(b) Γ_R *is planar if and only if* $p = 2$.

Proof. Note that $|Z(R)| = p$ and the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. So, by Theorem 2.3, we have

$$\text{Spec}(\Gamma_R) = \{(-1)^{p^3-2p-1}, (p^2-p-1)^{p+1}\} \text{ and } \gamma(\Gamma_R) = (p+1)\gamma(K_{p^2-p}).$$

Thus Γ_R is integral. Also, $\gamma(\Gamma_R) \neq 1$, that is, Γ_R is not toroidal. Part (b) follows from the fact that $\gamma(\Gamma_R) = 0$ if and only if $p = 2$. □

Proposition 3.4. *If* R *is a finite* 4*-centralizer ring then* Γ_R *is integral but not toroidal. Also* Γ_R *is planar if and only if* $|Z(R)| = 1, 2, 3$ *or* 4.

Proof. If R is a finite 4-centralizer ring then by [7, Theorem 3.2] we have that the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$. Now the results follow from Proposition 3.1. □

Proposition 3.5. *If R is a finite 5-centralizer ring then Γ_R is integral but not toroidal. Also Γ_R is planar if and only if $|Z(R)| = 1$ or 2.*

Proof. If R is a finite 5-centralizer ring then by [7, Theorem 4.3] we have that the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3$. Hence, the result follows from Proposition 3.1. □

In the following proposition, we compute the spectrum and genus of a $(p+2)$-centralizer p-ring, for any prime p.

Proposition 3.6. *If R is a finite $(p+2)$-centralizer p-ring, for any prime p, then*

$$\mathrm{Spec}(\Gamma_R) = \{(-1)^{(p^2-1)|Z(R)|-p-1}, ((p-1)|Z(R)|-1)^{p+1}\}$$

and $\gamma(\Gamma_R) = (p+1)\gamma(K_{(p-1)|Z(R)|})$.

Proof. If R is a finite $(p+2)$-centralizer p-ring then by [7, Theorem 2.12] the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$. Hence, the result follows from Theorem 2.3. □

The commuting probability of a ring R denoted by $\Pr(R)$ is the probability that a randomly chosen pair of elements of R commute. If R is a finite ring then $\Pr(R)$ is given by the ratio

$$\frac{|\{(r,s) \in R \times R : rs = sr\}|}{|R|^2}.$$

The study of $\Pr(R)$ was initiated by MacHale [13] in 1976. MacHale [13] proved the following result.

Theorem 3.7. *Let R be a finite ring and p the smallest prime divisor of $|R|$. Then $\Pr(R) \leq \frac{p^2+p-1}{p^3}$. The equality holds if and only if the additive quotient group $\frac{R}{Z(R)}$ is isomorphic to $\mathbb{Z}_p \times \mathbb{Z}_p$.*

As a consequence of Theorem 2.3 and Theorem 3.7 we have the following results.

Proposition 3.8. *Let R be a finite ring with $\Pr(R) = \frac{5}{8}$ then*

$$\mathrm{Spec}(\Gamma_R) = \{(-1)^{3(|Z(R)|-1)}, (|Z(R)|-1)^3\} \text{ and } \gamma(\Gamma_R) = 3\gamma(K_{|Z(R)|}).$$

Proposition 3.9. *Let R be a finite ring and p the smallest prime divisor of $|R|$. If $\Pr(R) = \frac{p^2+p-1}{p^3}$ then $\mathrm{Spec}(\Gamma_R) = \{(-1)^{(p^2-1)|Z(R)|-p-1}, ((p-1)|Z(R)|-1)^{p+1}\}$ and $\gamma(\Gamma_R) = (p+1)\gamma(K_{(p-1)|Z(R)|})$.*

References

[1] A. Abdollahi, Commuting graphs of full matrix rings over finite fields, *Linear Algebra Appl.* **428** (2008), 2947-2954.

[2] A. Abdollahi and E. Vatandoost, Which Cayley graphs are integral, *Electron. J. Combin.* **16**(2009), no. 1, R122, 17 pp.

[3] M. Afkhami, Z. Barati, N. Hoseini and K. Khashyarmanesh, A generalization of commuting graphs, *Discrete Math. Algorithm. Appl.* **7**(2015), no. 1, 1450068 (11 pages).

[4] S. Akbari, M. Ghandehari, M. Hadian and A. Mohammadian, On commuting graphs of semisimple rings, *Linear Algebra Appl.* **390** (2004), 345–355.

[5] S. Akbari and P. Raja, Commuting graphs of some subsets in simple rings, *Linear Algebra Appl.* **416** (2006), no. 23, 1038–1047.

[6] J. Battle, F. Harary, Y. Kodama and J. W. T. Youngs, Additivity of the genus of a graph, *Bull. Amer. Math. Soc.,* **68** (1962), 565–568.

[7] J. Dutta, D. K. Basnet and R. K. Nath, Characterizing some rings of finite order, submitted for publication, available at arXiv:1510.08207.

[8] J. Dutta, D. K. Basnet and R. K. Nath, On relative commuting graph of finite rings, preprint.

[9] J. Dutta and R. K. Nath, Finite groups whose commuting graphs are integral, submitted for publication.

[10] A. Erfanian, K. Khashyarmanesh and Kh. Nafar, Non-commuting graphs of rings, *Discrete Math. Algorithm. Appl.* DOI: 10.1142/S1793830915500275.

[11] F. Harary and A. J. Schwenk, *Which graphs have integral spectra?*, Graphs and Combinatorics, Lect. Notes Math., Vol 406, Springer-Verlag, Berlin, 1974, 45–51.

[12] G. Indulal and A. Vijayakumar, Some new integral graphs, *Appl. Anal. Discrete Math.* **1** (2007), 420–426.

[13] D. MacHale, Commutativity in finite rings, *Amer. Math. Monthly*, **83** (1976), 30–32.

[14] A. Mohammadian, On commuting graphs of finite matrix rings, *Comm. Algebra* **38** (2010), 988–994,

[15] R. K. Nath and J. Dutta, Spectrum of commuting graphs of some classes of finite groups, submitted for publication.

[16] G. R. Omidi and E. Vatandoost, On the commuting graph of rings, *J. Algebra Appl.* **10** (2011), no. 3, 521–527.

[17] E. Vatandoost, F. Ramezani and A. Bahraini, On the commuting graph of non-commutative rings of order $p^n q$, *J. Linear Topological Algebra* **3** (2014), no. 1, 1–6.

[18] L. Wang, X. Li and C. Hoede, Two classes of integral regular graphs, *Ars Combin.* **76** (2005), 303–319.

[19] A. T. White, Graphs, Groups and Surfaces, *North-Holland Mathematics Studies, no. 8.,* American Elsevier Publishing Co., Inc., New Yourk, 1973.